

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/351287551>

Vitamin D Deficiency in Brazil: Questions about Potential Causes and Clinical Characteristics

Article in *International Journal of Nutrology* · May 2021

DOI: 10.11055/s-0041-1728682

CITATIONS

0

READS

61

6 authors, including:



Thiago Hirose

Centro Universitário Estácio de Ribeirão Preto

5 PUBLICATIONS 0 CITATIONS

SEE PROFILE



Renato Augusto Zorzo

Universidade Estácio de Sá

9 PUBLICATIONS 11 CITATIONS

SEE PROFILE



Vitamin D Deficiency in Brazil: Questions about Potential Causes and Clinical Characteristics*

Deficiência de vitamina D no Brasil: Questionamentos sobre as suas possíveis causas e características clínicas

Fabília Belloni dos Santos Vieira¹ Ana Paula Morguete Ferreira¹ Felipe Rezende Giacomelli¹
Lívia Gonçalves de Lima¹ Thiago Santos Hirose¹ Renato Augusto Zorzo²

¹ Centro Universitário Estácio de Ribeirão Preto, Ribeirânia, Ribeirão Preto, SP, Brazil

² Universidade Federal de São Carlos, São Carlos, SP, Brazil

Address for correspondence Lívia Gonçalves de Lima, Centro Universitário Estácio de Ribeirão Preto, – Rua Abrahão Issa Halack, 980, Ribeirânia, Ribeirão Preto, SP, 14096-160, Brazil (e-mail: liviagli@outlook.com).

Int J Nutrol 2021;14:26–32.

Abstract

Vitamin D is a micronutrient essential to various systems from the human body, and it is not restricted to the classical function of bone mineralization. Its synthesis is mainly related to ultraviolet B (UVB) radiation exposure. Although Brazil is a tropical country with high levels of UVB radiation, counter-intuitively, a large number of Brazilians present vitamin D deficiency, which is also a worldwide issue. This review aims to approach clinical features and explore potential causes for this apparent contradiction through questions that could explain vitamin D deficiency in the Brazilian population.

Keywords

- ▶ Vitamin D
- ▶ Avitaminosis
- ▶ Ultraviolet Rays

Resumo

A vitamina D corresponde a um micronutriente essencial ao funcionamento de diversos sistemas do corpo humano, não se restringindo à clássica função de mineralização óssea. A sua síntese está relacionada principalmente à exposição à radiação ultravioleta B (UVB). O Brasil é um país tropical que apresenta altos índices de UVB, no entanto, ao contrário do que se pode imaginar, observa-se um grande número de brasileiros deficientes em vitamina D e essa deficiência se torna um problema de ordem mundial. Essa revisão tem por objetivo abordar características clínicas e explorar as possíveis causas para essa aparente contradição, por meio de questionamentos que poderiam explicar a deficiência de vitamina D na população brasileira.

Palavras-chave

- ▶ Vitamina D
- ▶ Deficiência de Vitaminas
- ▶ Raios Ultravioleta

Introduction

Vitamin D is a fat-soluble steroid hormone, and it is considered a micronutrient essential to a complex endocrinological

system.^{1–6} Its action has been studied since the beginning of the 20th century and it is increasingly clear that vitamin D functions are not restricted to the classical bone mineralization regulation.^{1–3,7}

Vitamin D has several roles, encompassing both endocrine reactions and functions from other systems, such as the immune system. Its protective immunomodulatory role in

* Study performed at Centro Universitário Estácio de Ribeirão Preto, Ribeirão Preto, SP, Brazil.

received
December 11, 2020
accepted
February 3, 2021

DOI <https://doi.org/10.1055/s-0041-1728682>.
ISSN 1984-3011.

© 2021. Associação Brasileira de Nutrologia. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil




Fig. 1 Vitamin D metabolism. Source: Modified from "Alimentação, fotoexposição e suplementação: influência nos níveis séricos de vitamina D" (2015) and "Sunlight exposure: Do health benefits outweigh harm?" (2016).

conditions such as systemic arterial hypertension, cancer, osteoarthritis, multiple sclerosis, and diabetes mellitus is described.^{1,2,8}

Although the main site of vitamin D metabolism (► **Figure 1**) is the skin, it also occurs in other properly functioning organs, such as the bowel, liver and kidney.^{9,10} Cutaneous 7-dehydrocholesterol (pro-vitamin D₃) is converted by ultraviolet B (UVB) radiation in cholecalciferol (vitamin D₃), which can be also obtained from food sources.^{9,10} Within the liver, an enzyme, called 25-hydroxylase, converts vitamin D₃ in 25-hydroxy vitamin D₃ (25 (OH) vitamin D) or calcidiol.^{9,10} Calcidiol passes through the kidneys and undergoes a second hydroxylation by an enzyme called 1-alpha-hydroxylase, resulting on 1,25-dihydroxy-vitamin D₃ (calcitriol), the active form of the molecule.^{9,10} This enzyme is activated by parathyroid hormone and calcium and phosphate concentrations, demonstrating the close relationship between vitamin D and bone metabolism.^{9,10}

Skin exposure to UVB radiation is the main source of vitamin D.^{7,11,12} Vitamin D can be obtained from food, but with a lower efficiency compared to sunlight.^{13,14} Vitamin D deficiency is a global health issue.^{4,15} At first glance, it can be attributed to a reduction in UVB exposure, especially in countries with lower solar incidence.

However, tropical countries, such as Brazil, have a high UVB incidence and, despite the lack of populational studies, number of people with vitamin D deficiency is increasing.^{16,17} This is curious given that UVB is the main source of vitamin D for the human body.^{7,12,13}

Review

Data to effectively answer these questions were obtained from 57 sources describing the situation in Brazil, the United States, India, and France, showing that vitamin D, popularly known as the "sunshine vitamin,"¹⁸ is extremely important and derived in different ways. UVB is the primary source of endogenously synthesized vitamin D, representing 80 to 90% of the daily fraction required by the body.^{19,20} The spectrum of UVB radiation considered optimal for pre-vitamin D synthesis at the skin ranges from 290 to 320 nm; coincidentally, this is the same range associated with skin carcinogenesis.²¹⁻²³ In Brazil, the average daily UVB radiation ranges from 3,300 to 3,800 J/m², reaching values greater than 7,000 J/m² during the summer.^{21,24} For people with light skin, very sensitive to the sun and with little pigmentation (phototype II), the exposure of small, uncovered skin areas for about 10 minutes is sufficient to absorb the required amount of UVB for vitamin D synthesis. In contrast, people with darker skin (phototype IV) would need a longer exposure time, since melanin absorbs UV radiation and limits the number of photons available for pre-vitamin D₃ formation.²⁴

Vitamin D-rich foods constitute a secondary source for this nutrient, accounting for 10% to 20% of the body requirements.¹⁹ Therefore, vitamin D is rarely obtained in sufficient quantities with common diets.²⁵ The main sources of vitamin D₃ are fatty fish from cold, deep waters, such as salmon and tuna.¹⁹ Soy milk, margarine, breads, breakfast cereals, orange juice, and egg yolks are also good vitamin D sources.^{1,10,26-28}

Brazil has a difficult ethnic classification due to its high degree of miscegenation.²⁹ As revealed by the 2016 National Continuous Household Sample Survey (PNAD), the Brazilian population is comprised of 8.2% black and 46.2% brown subjects. Thus, most Brazilians present higher phototypes, which could corroborate for the perceived vitamin D deficiency.

This hypovitaminosis usually occurs in obese individuals in a 50% higher proportion compared to eutrophic subjects.³⁰ This finding may be related to the presence of vitamin D receptors in the adipose tissue, which would extract the vitamin from the blood circulation, reducing its concentration. In addition, obesity tends to reduce sunlight exposure due to mobility limitations.^{31,32}

It is estimated that approximately 80% of the world population live in areas with air pollution levels higher than those defined by air quality guidelines.^{33,34} In Brazil, 75% of the population live in urban centers, where air pollution is a reality, especially in large cities. Thus, people from large Brazilian industrial centers potentially present higher rates of vitamin D deficiency.¹¹

Serum vitamin D level is assessed by measuring the fraction of 25 (OH) vitamin D using mass spectrometry (considered the golden standard method).⁷ This choice is based on the fact that calcidiol is the main metabolite in the bloodstream when compared to the active form of the vitamin. Vitamin D deficiency, insufficiency, and sufficiency are based on 25 (OH) vitamin D levels (► **Table 1**).³⁵⁻³⁸

Table 1 Reference values for 25 (OH) vitamin D according to the US Endocrine Society

Definition	Reference Values*
Sufficiency	30 to 100 ng/mL
Insufficiency	20 to 29 ng/mL
Deficiency	Lower than 20 ng/mL

*Reference values for 25 (OH) vitamin D

Source: Adapted from "Recomendações da Sociedade Brasileira de Endocrinologia e Metabologia (SBEM) para o diagnóstico e tratamento de hipovitaminose D" (2014).

What are the main risk groups for vitamin D deficiency?

Premature children; newborns exclusively receiving breast milk, which has a low amount of vitamin D; obese subjects; elderly people; pregnant women; patients with rickets/osteomalacia and osteoporosis; patients with a history of fall and fracture due to a deficiency which impairs muscle relaxation and contraction, resulting in pain and weakness;³⁹ subjects with hyperparathyroidism; inflammatory diseases; autoimmune diseases, such as multiple sclerosis, asthma and rheumatoid arthritis;^{40,41} chronic kidney disease; and malabsorption syndromes.⁴²

What is the clinical picture of vitamin D deficiency?

When vitamin D deficiency is not asymptomatic, patients may report irritability, sweating and even more severe conditions, such as rickets in children and osteomalacia in adolescents and adults. These more severe conditions result from hypocalcemia and hypophosphatemia, respectively caused by a reduced calcium and phosphate absorption and an increased phosphate clearance at the intestinal level.³⁹

Are Brazilians not exposed to sunlight enough?

Lifestyle changed in recent years, especially regarding sunlight exposure. During the last century, Brazil has undergone an intense process of urbanization and economic development. An essentially rural population started to concentrate in cities. In urban centers, work is carried out basically indoors. There was also a reduction in outdoor physical activity and an increase in sedentary lifestyle, resulting in obesity.⁴³⁻⁴⁶

In addition, sunscreen use has become widespread. At first, it was assumed that sunscreens were responsible for vitamin D levels reduction both in the Brazilian and global populations.²⁴ Indeed, these products are effective in preventing UVB penetration into the skin. However, it is worth mentioning that sunscreens only provide the UVB protection indicated on the label when applied in an amount of 2 mg/cm² every two hours.²⁴ Routinely, few people use sunscreens properly to be fully protected from UVB radiation.

Data show that sunlight exposure of Brazilian people was reduced, either due to socioeconomic changes or to the

increased use of sunscreens. However, the incidence of UVB in the country is high and the amount of UVB required for vitamin D synthesis is low, which calls into question this hypothesis as the only cause for the hypovitaminosis D seen in Brazil.

Is the Brazilian diet low in vitamin D?

The daily requirement for vitamin D is 1,500 to 2,000 IU/day (37.5 to 50 µg) for healthy adults.¹⁰ It is unlikely that this dose is obtained from diet alone, even when foods with a very high content of vitamin D, such as those shown in **Table 2**, are consumed.

It is difficult to obtain enough vitamin D only from food in Brazil, since the basic local diet consists of items with a high nutritional content (rice, beans, milk, meat, and fruits), but with a low vitamin D index.⁴⁷ Thus, although the Brazilian diet can contribute to this hypovitaminosis, it is not its main cause, since foods are secondary sources.

Could Brazilian population miscegenation result in a reduced vitamin D synthesis?

Skin pigmentation affects the cutaneous synthesis of vitamin D. Therefore, people with a higher concentration of melanin present lower vitamin D synthesis because the pigment acts as a natural sunscreen, reducing UVB absorption.^{16,46}

However, it is worth mentioning that such data are self-reported. In addition, there is a lack of studies linking miscegenation to vitamin D deficiency.

Could the increase in obesity in the Brazilian population interfere with vitamin D levels?

Obesity is a current worldwide public health problem. Data from the World Health Organization (WHO) revealed that, in 2016, more than 1.9 billion adults, aged 18 years old or over, were overweight, including more than 650 million obese subjects.⁴⁸ In Brazil, the situation is similar, and it is

Table 2 Main vitamin D-rich foods: note that few are commonly included in the Brazilian diet

Food	Portion size	Vitamin D (UI)
Fish liver oil	23.1 mg/ 1 tablespoon	924
Grilled salmon	100 g	284
Grilled mackerel	100 g	352
Canned tuna in brine	100 g	144
Canned sardine in brine	100 g	184
Chicken egg	50g/ 1 regular unit	3
Fried beef liver	100g	36
Fortified margarine	20g	62
Fortified breakfast cereal	30g/medium-sized portion	52

Source: Modified from "Funções Plenamente Reconhecidas de Nutrientes Vitamina D" (2015).

estimated that 49% of the population is overweight and 14.8% is obese according to studies from the Health Department.⁴⁹

Dyslipidemia is often associated with obesity. According to the 2013 National Health Survey (PNS), 12.5% of the Brazilian population aged 18 and over (18.4 million people) present dyslipidemia.⁵⁰ Dyslipidemia patients are instructed to not eat foods that are good vitamin D sources, including egg yolk, liver, butter, and milk, since they present a high fat content.⁵¹

Therefore, the higher obesity rates in the Brazilian population are potentially related to the increase in perceived hypovitaminosis D. There are no further studies on this relationship, especially in Brazil.

Does pollution in large Brazilian centers affect vitamin D synthesis?

There are no relevant studies associating air pollution levels to hypovitaminosis D. However, it is known that air-borne impurities can absorb part of the UVB radiation, decreasing the amount reaching Earth's surface. Thus, more polluted areas may have less UVB radiation, contribution to a reduction in vitamin D levels.^{33,52,53}

Studies from India and France debate the relationship between vitamin D deficiency and air pollution. A study carried out with French mothers demonstrated a potential correlation between low levels of 25 (OH) D at birth and the mother's exposure to air pollution during pregnancy.⁵² The Indian study (evaluating vitamin D status in babies and children living in similar house conditions, but in two city areas with different air pollution levels) revealed significantly lower levels of vitamin D in subjects from the polluted area in comparison to those from the less polluted area.⁵⁴

It is also questioned whether, in contrast, subjects from rural areas would have higher vitamin D levels compared to city dwellers.

Are vitamin D reference values correct for the Brazilian population?

There is a worldwide consensus that serum calcidiol levels below 25 to 30 nmol/L can damage bone metabolism.^{1,55} Within this context, it is worth highlighting a discussion regarding these values, which change according to location and source, as shown in ►Table 1.

Currently, most services use the values suggested by the US Endocrine Society. However, their determination was based on a population sample, whose characteristics may not be representative of a given country's population.

As such, it is possible that the adopted reference levels are not suitable for the Brazilian population due to its sociocultural and environmental peculiarities.^{55,56} Further studies are required to assess normal serum levels of vitamin D considering Brazilian particularities.

May values change according to sample collection, habits, and characteristics of a country's population?

According to the habits and behavior of the Brazilian population, sunscreen use may be associated with lowest 25 (OH)

vitamin D levels in comparison with sunlight-exposed subjects.²¹

In addition, genetic features play a role, since it is known that the phosphate-regulating gene with homologies to endopeptidases on the X chromosome (PEX) alters the function of one of the proteins acting on phosphate regulation, which is closely linked to the diagnosis of rickets due to hypovitaminosis D.⁵⁷

Therefore, it is essential to consider behaviors and even clinical issues from each patient to determine whether there is a diagnosis of vitamin deficiency or if these are only factors inducing the condition.

Conclusion

In a country with high UVB levels, such as Brazil, it is curious to talk about vitamin D deficiency, since UVB is the main source for the synthesis of the popular "sunshine vitamin". A few minutes under the Brazilian sun would be enough to obtain the required UVB dose for daily vitamin D synthesis.

However, a large portion of the Brazilian population presents vitamin D deficiency, although there are no population studies quantifying it. Therefore, its potential causes are questioned.

The hypotheses raised and explored in this review indicate that reduced vitamin D levels do not result from a single factor, but a combination of sociocultural, environmental, and physiological factors, along with divergences regarding the adopted reference values.

However, there are no specific studies on the Brazilian population to elucidate the real causes of vitamin D deficiency. The questions addressed here have no definitive answers, but they would be good starting points for new inquiries.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- Gallego-González D, Mejia-Mesa S, Martínez-Sánchez LM, Rendón-Diez M. Hipovitaminosis D: una visión desde la clínica y la biología molecular. *Med UIS*. 2017;30(01):45–56. Doi: 10.18273/revmed.v30n1-2017004
- Chen S, Sun Y, Agrawal DK. Vitamin D deficiency and essential hypertension. *J Am Soc Hypertens* 2015;9(11):885–901. Doi: 10.1016/j.jash.2015.08.009
- Hilger J, Friedel A, Herr R, et al. A systematic review of vitamin D status in populations worldwide. *Br J Nutr* 2014;111(01):23–45. Doi: 10.1017/S0007114513001840
- Ferrer CF, Mercadé MVF. Deficiência de vitamina D (25OH) e seu impacto na qualidade de vida. *RBAC* 2015;32(02):77–79. Doi: 10.21877/2448-3877.201800686
- IARC Working Group on Vitamin D. Vitamin D and cancer: a report of the IARC Working Group on Vitamin D. Lyon: IARC; 2008(IARC Working Group Reports; 5). <https://doi.org/10.4161/derm.1.1.7729>
- Rosen CJ, Adams JS, Bikle DD, et al. The nonskeletal effects of vitamin D: an Endocrine Society scientific statement. *Endocr Rev* 2012;33(03):456–492. Doi: 10.1210/er.2012-1000
- da Silva Quadros KR, de Oliveira RB. Reposição de vitamina D nativa: indicação à luz das evidências científicas atuais. *Rev Fac*

- Ciênc Méd Sorocaba. 2016;18(02):79–86. Doi: 10.5327/z1984-4840201626833
- 8 Berridge MJ. Vitamin D: a custodian of cell signalling stability in health and disease. *Biochem Soc Trans* 2015;43(03):349–358. Doi: 10.1042/BST20140279
 - 9 Grüttnert VS, Weingrill P, Fernandes AL. Absorption aspects of calcium and vitamin D metabolism. *Rev Bras Reumatol* 1997;37(03):143–151 Disponível em: <http://bases.bireme.br/cgi-bin/wxislind.exe/iah/online/?IscScript=iah/iah.xis&src=google&base=LILACS&lang=p&nextAction=lnk&exprSearch=210140&indexSearch=ID#refine>
 - 10 Emo B, Araújo L. Funções plenamente reconhecidas de nutrientes vitamina D. *ILSI Brasil* 2015;2:3–18. Disponível em: https://ilsio.org/brasil/wp-content/uploads/sites/9/2016/05/artigo_vitamina_D_espanhol-FINAL.pdf
 - 11 Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357(03):266–281. Doi: 10.1056/NEJMra070553
 - 12 Holick MF, Garabedian M. Vitamin D: photobiology, metabolism, mechanism of action, and clinical application. In: Favus MJ, editor. *Primer on the metabolic bone diseases and disorders of mineral metabolism*. 6th ed. Pennsylvania: Rittenhouse Book Distribution; 2006:106–14. Available from: <https://opac.ll.chiba-u.jp/da/curator/900030218/>
 - 13 Marques CDL, Dantas AT, Fragoso TS, Duarte ALBP. A importância dos níveis de vitamina D nas doenças autoimunes. *Rev Bras Reumatol* 2010;50(01):67–80. Doi: 10.1590/S0482-5004201000100007
 - 14 Bringham FR, Demay MB, Kronenberg HM. Hormones and Disorders of Mineral Metabolism. In: Kronenberg HM, Melmed S, Polonsky KS, Larsen PR, editors. *Williams textbook of endocrinology*. 11th ed. Philadelphia: Elsevier; 2008
 - 15 Palacios C, Gonzalez L. Is vitamin D deficiency a major global public health problem? *J Steroid Biochem Mol Biol* 2014;144(Pt A):138–145. Doi: 10.1016/j.jsbmb.2013.11.003
 - 16 de Carvalho Mesquita K, de Souza Machado Igreja AC, Costa IMC. Dermatite atópica e vitamina D: Fatos e controvérsias. *An Bras Dermatol* 2013;88(06):945–953. Doi: 10.1590/abd1806-4841.20132660
 - 17 Peters BS, dos Santos LC, Fisberg M, Wood RJ, Martini LA. Prevalence of vitamin D insufficiency in Brazilian adolescents. *Ann Nutr Metab* 2009;54(01):15–21. Doi: 10.1159/000199454
 - 18 Holick MF. Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am J Clin Nutr* 2004;80(6, Suppl):1678S–1688S. Doi: 10.1093/ajcn/80.6.1678S
 - 19 de Castro LCG. O sistema endocrinológico vitamina D. *Arq Bras Endocrinol Metab* 2011;55(08):566–575. Doi: 10.1590/S0004-27302011000800010
 - 20 Holick MF. Vitamin D: a D-Lightful health perspective. *Nutr Rev* 2008;66(10, Suppl 2):S182–S194. Doi: 10.1111/j.1753-4887.2008.00104.x
 - 21 Maia M, Maeda SS, Marçon C. Correlação entre fotoproteção e concentrações de 25 hidróxi-vitamina D e paratormônio. *An Bras Dermatol* 2007;0682(03):233–237. Doi: 10.1590/S0365-05962007000300004
 - 22 Cole CA, Forbes PD, Davies RE. An action spectrum for UV photocarcinogenesis. *Photochem Photobiol* 1986;43(03):275–284. Doi: 10.1111/j.1751-1097.1986.tb05605.x
 - 23 Holick MF. The photobiology of vitamin D and its consequences for humans. *Ann N Y Acad Sci* 1985;453:1–13. Doi: 10.1111/j.1749-6632.1985.tb11793.x
 - 24 Schalka S, Steiner D, Ravelli FN, et al. Consenso Brasileiro de Fotoproteção. *An Bras Dermatol* 2014;89(6 Suppl 1):56–75 <http://www.sbd.org.br/publicacoes/consenso-brasileiro-de-fotoprotecao>
 - 25 Arnson Y, Amital D, Amital H. The diverse world of vitamin D: does it also modulate pain sensation? *Isr Med Assoc J* 2009;11(06):371–372 <https://pubmed.ncbi.nlm.nih.gov/19697590/>
 - 26 Wacker M, Holick MF. Sunlight and Vitamin D: A global perspective for health. *Dermatoendocrinol* 2013;5(01):51–108. Doi: 10.4161/derm.24494
 - 27 Holick MF. Optimal vitamin D status for the prevention and treatment of osteoporosis. *Drugs Aging* 2007;24(12):1017–1029. Doi: 10.2165/00002512-200724120-00005
 - 28 Kennel KA, Drake MT, Hurley DL. Vitamin D. Vitamin D deficiency in adults: when to test and how to treat. *Mayo Clin Proc* 2010;85(08):752–757, quiz 757–758. Doi: 10.4065/mcp.2010.0138
 - 29 Damaso ÊL, Paula FJ, Franceschini SA, et al. Does the Access to Sun Exposure Ensure Adequate Levels of 25-Hydroxyvitamin D? *Rev Bras Ginecol Obstet* 2017;39(03):102–109. Doi: 10.1055/s-0037-1600520
 - 30 Medeiros M, Saunders C, Chagas CB, Pereira SE, Saboya C, Ramalho A. Vitamin D deficiency in pregnancy after bariatric surgery. *Obes Surg* 2013;23(10):1679–1684. Doi: 10.1007/s11695-013-1045-5
 - 31 Silva JDS, Pereira S, Saboya Sobrinho C, Ramalho A. Obesity, related diseases and their relationship with vitamin D deficiency in adolescents. *Nutr Hosp* 2016;33(04):856–864. Doi: 10.20960/nh.381
 - 32 Kull M, Kallikorm R, Lember M. Body mass index determines sunbathing habits: implications on vitamin D levels. *Intern Med J* 2009;39(04):256–258. Doi: 10.1111/j.1445-5994.2009.01900.x
 - 33 Barrea L, Savastano S, Di Somma C, et al. Low serum vitamin D status, air pollution and obesity: A dangerous liaison. *Rev Endocr Metab Disord* 2017;18(02):207–214. Doi: 10.1007/s11154-016-9388-6
 - 34 van Donkelaar A, Martin RV, Brauer M, et al. Global estimates of ambient fine particulate matter concentrations from satellite-based aerosol optical depth: development and application. *Environ Health Perspect* 2010;118(06):847–855. Doi: 10.1289/ehp.0901623
 - 35 Maeda SS, Borba VZC, Camargo MBR, et al. Recomendações da Sociedade Brasileira de Endocrinologia e Metabologia (SBEM) para o diagnóstico e tratamento de hipovitaminose D. *Arq Bras Endocrinol Metab* 2014;58(05):411–433. Doi: 10.1590/0004-27300000003388
 - 36 Norman AW, Bouillon R. Vitamin D nutritional policy needs a vision for the future. *Exp Biol Med (Maywood)* 2010;235(09):1034–1045
 - 37 Holick MF, Binkley NC, Bischoff-Ferrari HA, et al; Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2011;96(07):1911–1930. Doi: 10.1210/jc.2011-0385
 - 38 Heaney RP. What is vitamin D insufficiency? And does it matter?. *Calcif Tissue Int* 2013;92(02):177–183. Doi: 10.1007/s00223-012-9605-6
 - 39 Premaor MOE, Furlanetto TW. Hipovitaminose D em adultos: entendendo melhor a apresentação de uma velha doença. *Arq Bras Endocrinol Metab* 2006;50(01):25–37. Doi: 10.1590/S0004-27302006000100005
 - 40 Dobnig H, Pilz S, Scharnagl H, et al. Independent association of low serum 25-hydroxyvitamin d and 1,25-dihydroxyvitamin d levels with all-cause and cardiovascular mortality. *Arch Intern Med* 2008;168(12):1340–1349. Doi: 10.1001/archinte.168.12.1340
 - 41 Souberbielle J-C, Body JJ, Lappe JM, et al. Vitamin D and musculoskeletal health, cardiovascular disease, autoimmunity and cancer: Recommendations for clinical practice. *Autoimmun Rev* 2010;9(11):709–715. Doi: 10.1016/j.autrev.2010.06.009
 - 42 Ferreira CES, Maeda SS, Batista MC, et al. Consensus – reference ranges of vitamin D [25(OH)D] from the Brazilian medical societies. Brazilian Society of Clinical Pathology/Laboratory Medicine (SBPC/ML) and Brazilian Society of Endocrinology and Metabolism (SBEM). *J Bras Patol Med Lab* 2017;53(06):377–381. Doi: 10.5935/1676-2444.20170060
 - 43 Moy FM, Hoe VCNN, Hair NN, Vethakkan SR, Bulgiba A. Vitamin D deficiency and depression among women from an urban

- community in a tropical country. *Public Health Nutr* 2017;20(10): 1844–1850. Doi: 10.1017/S1368980016000811
- 44 Ginde AA, Liu MC, Camargo CA Jr. Demographic differences and trends of vitamin D insufficiency in the US population, 1988–2004. *Arch Intern Med* 2009;169(06):626–632. Doi: 10.1001/archinternmed.2008.604
- 45 Holick MF. Resurrection of vitamin D deficiency and rickets. *J Clin Invest* 2006;116(08):2062–2072. Doi: 10.1172/JCI29449
- 46 Scragg R, Camargo CA Jr. Frequency of leisure-time physical activity and serum 25-hydroxyvitamin D levels in the US population: results from the Third National Health and Nutrition Examination Survey. *Am J Epidemiol* 2008;168(06):577–586, discussion 587–591. Doi: 10.1093/aje/kwn163
- 47 Pereda PC, de Oliveira Alves DC. Qualidade alimentar dos brasileiros: teoria e evidência usando demanda por nutrientes. *Pesqui Planej Econ* 2012;42(02):239–260 Disponível em: <http://repositorio.ipea.gov.br/handle/11058/4960>
- 48 World Health Organization. Obesity and Overweight. Available from: <https://www.who.int/westernpacific/health-topics/obesity>
- 49 Macari C, et al. Obesidade, perfil lipídico e hábitos alimentares de escolares: comparação entre municípios de dois estados da região sul do Brasil. *Rev Saúde e Pesqui.* 2017;10(03):451–461. Doi: 10.17765/2176-9206.2017v10n3p451-461
- 50 Instituto Brasileiro de Geografia e Estatística – IBGE. Pesquisa Nacional de Saúde. Disponível em <https://www.ibge.gov.br/estatisticas-novoportal/sociais/saude/9160-pesquisa-nacional-de-saude.html?=&t=o-que-e>
- 51 de Oliveira Andrade PC, de Castro LS, de Souza Lambertucci M, et al. Diet, sun exposure, and dietary supplementation: effect on sérum levels of vitamin D. *Rev Med Minas Gerais.* 2015;25(03): 432–437. Doi: 10.5935/2238-3182.20150082
- 52 Baiz N, Dargent-Molina P, Wark JD, Souberbielle JC, Slama R, Annesi-Maesano IEDEN Mother-Child Cohort Study Group. Gestational exposure to urban air pollution related to a decrease in cord blood vitamin d levels. *J Clin Endocrinol Metab* 2012;97(11): 4087–4095. Doi: 10.1210/jc.2012-1943
- 53 Lange NE, Litonjua A, Hawrylowicz CM, Weiss S. Vitamin D, the immune system and asthma. *Expert Rev Clin Immunol* 2009;5 (06):693–702. Doi: 10.1586/eci.09.53
- 54 Agarwal KS, Mughal MZ, Upadhyay P, Berry JL, Mawer EB, Puliyl JM. The impact of atmospheric pollution on vitamin D status of infants and toddlers in Delhi, India. *Arch Dis Child* 2002;87(02): 111–113. Doi: 10.1136/adc.87.2.111
- 55 Thienpont LM, Stepman HC, Vesper HW. Standardization of measurements of 25-hydroxyvitamin D3 and D2. *Scand J Clin Lab Invest Suppl* 2012;243:41–49. Doi: 10.3109/00365513.2012.681950
- 56 Torrubia B, Alonso I, Mahillo I, et al. Comparación entre dos inmuno-ensayos automatizados por quimio-luminiscencia para la cuantificación de 25 (OH) vitamina D. *Rev Osteoporos Metab Miner.* 2016;8(02):70–74 Disponível em: http://scielo.isciii.es/scielo.php?script=sci_arttext&pid=S1889-836X22016000200004&lng=es
- 57 Nunes AB, Lazaretti-Castro M. Raquitismo hipofosfatêmico: da clínica à genética molecular. *Arq Bras Endocrinol Metabol* 2000; 44(02):125–132. Doi: 10.1590/S0004-2730200000200004